

Sw., *Ephemerantha* P.F.Hunt & Summerh., *Eria* Lindl., *Galeola* Lour., *Gastrodia* R.Br., *Gymnadenia* R.Br., *Habenaria* Willd., *Ludisia* A.Rich., *Luisia* Gaudich., *Nervilia* Comm. ex Gaudich., and *Thunia* Rchb.f. are reported as the frequently used genera in medicine. Several bioactive compounds such as alkaloids (dendrobine, jebantine), bibenzyl (gigantol, moscatilin), phenanthrene (nudol, agrostophyllinone, isoagrostophyllol, nidemin), phenanthrenequinone (cypripedin), glycosides (loroglossin, melianin, kinsenoside), carbohydrates, flavonoids, stilbenoids (arundinan, orchinol, 6-methoxycoelonin, imbricatin, flaccidin, oxoflaccidin, isooxoflaccidin, flaccidin, agrostophyllin, callosin, callosinin, callosumin, callosuminin and callosumidin) have been reported from several orchid species²⁰⁻²³. These orchid compounds are useful in healing fractures, seminal weakness, fever, abnormal thirst, curing cancer, and diabetic conditions²³⁻²⁴. Several bioactive compounds found in orchids have been shown to possess various pharmacological properties such as anti-microbial, anti-inflammatory, anti-tumour, and anti-viral²⁴. Due to human activities, these groups of plants have a higher proportion of threatened and endangered species.

Keeping in view the importance of traditional knowledge associated with plants in drug discovery programs, an investigative study was carried out to document the medicinal orchids of the Himalaya regions used by nomadic people.

Methodology

Study area

Indian Himalaya (26°20' to 35°40'N, 74°50' to 95°40'E) with 250 to 300 km is spread over 2,500 km stretch from union territory Jammu & Kashmir (J&K) in the west to Arunachal Pradesh in the east^{6,18}. It borders Pakistan on the west, Myanmar in the Eastern Himalaya, touching parts of Nepal and Bhutan. The region partially/full covers 11 states Himachal Pradesh, Uttaranchal, Sikkim, Arunachal Pradesh, Nagaland, Manipur, Mizoram, Tripura, Meghalaya, Assam, West Bengal, and 1 union territory J&K. The region has a total geographical area of approximately 5,33,604 km² and is home to 3,69,28,311 people¹⁸. The region supports unique cultural, ethnic, and biological diversity and there are several glaciers such as Siachen, Rulung, Neh-Nar, Braham Sar, and Harmukh originating from the high hills mountains of the Indian Himalaya¹⁸. Climate is characterized by tropical/subtropical climatic

conditions from the foothill region of the mountains to permanent ice and snow-covered peaks at higher altitudes^{11,12}. The species of plants and animals of the Himalayas vary with climate, rainfall, altitude, and soils. The amount of annual rainfall increases from west to east along the southern front of the range. The valleys experience a mean summer temperature between 15 and 25 °C and a much colder winter; the north-western peaks of Himalaya typically experience dry conditions, with surface temperatures ranging between 3 and 35 °C in summer and -20 and -35 °C in winter together with heavy snowfall^{16,18}. The mountains have rich forests abode with thousand of species of trees, shrubs, herbs, lianas and more than 40% of the species are endemics. Many medicinal species are growing in the Himalayas which have high demand in the international trade such as species of *Aconitum* L., *Allium* L., *Angelica* L., *Cinnamomum* Schaeff., *Panax* L., *Paris* L., *Podophyllum* L., *Saussurea* DC., *Terminalia* L., and *Zanthoxylum* L.^{13,18}.

Field survey and ethnobotanical documentation

Field surveys, investigations, and plant collection tours for studying medicinal orchids were conducted between 2015 and 2018. Few pockets of Himalaya regions were selected based on their known cultural and botanical richness. Before the field-works, relevant information and literature²⁵⁻³⁵ were reviewed with reference to local indigenous knowledge on people, topography, and climate of the study areas. Field studies were undertaken during the growing season by interviewing members of the local nomadic communities inhabiting the Indian Himalayas. Medicinal information on orchid plants was collected from 146 individuals (81 males and 65 females) (Table 1). The data collected were from headmen, heads of the family, and herbal practitioners, through direct observation, and individual discussions. The plant materials collected as voucher samples were prepared in the usual way³⁶ and stored in the ASSAM herbarium. The acronym of the herbarium names is in accordance with Thiers³⁷.

Plant authentication and certification

All local plant names recorded during field were scientifically authenticated using keys and descriptions from literature (regional floras, manuals, monographs), research articles and by matching the collected samples with that of deposited vouchers at herbaria of the Botanical Survey of India, Eastern Regional Centre, Shillong; The Herbarium of Forest

Table 1 — Surveyed area and informants involved in documentation of traditional knowledge of Himalayan orchids

Surveyed Area	Year of survey	Sex and number of informants			GPS Coordinates		
		Male	Female	Total	Latitude (N)	Longitude (E)	Altitude (m amsl)
Jammu & Kashmir	2015 (July, September)	9	4	13	34°38'14.06"	74°52'07.04"	2692
Himachal Pradesh	2012 (April)	10	1	11	32°03'42.12"	77°15'40.68"	3029
Uttarakhand	2011 (June)	7	1	8	30°05'12.84"	78°16'03.36"	2189
Sikkim	2010 (August)	3	14	17	27°37'57.07"	88°45'15.01"	2700
Arunachal Pradesh	2013 (September)	13	16	29	27°15'53.00"	92°25'28.00"	2507
Meghalaya	2011 (March)	18	20	38	25°31'60.01"	90°06'60.08"	1050
Assam	2011 (January)	21	9	30	26°34'39.00"	93°10'15.96"	354
Total		81	65	146			

Research Institute, Dehradun; Janaki Ammal Herbarium, Jammu; Gauhati University, Guwahati for Indian Himalayan species. Valid botanical names along with the author citations were verified from Plants of the World Online (<http://www.plantsoftheworldonline.org/>). All vouchers were deposited in ASSAM herbarium at Botanical Survey of India, Eastern Regional Centre Shillong.

Data analysis

Informed consent was obtained verbally from all the informants prior to the semi-structured interviews. The data collected through discussions and interviews while surveying the localities were analyzed using quantitative index use frequency (F_q), which is expressed in percentage and calculated by multiplying the use-value mentioned by informants with 100. This gives the overall percentage of use reports of the particular species of the study area.

Results and Discussion

Population structure of informants

Table 1 shows the area surveyed, period of field trips, number of informants, and GPS coordinates recorded during the investigations of the traditional knowledge associated with orchids growing in the Himalaya regions. The Meghalaya state recorded the largest number of informants (38) having the indigenous knowledge on the use of orchid species as medicine, followed by informants of Assam (30) and Arunachal Pradesh (29). The state of Uttarakhand recorded the smallest number of informants (8). In total, 146 informants (81 males and 65 females) were interviewed from the different study sites. Respondents belong to all age groups (I: 15-30 yrs, II: 31-45 yrs, III: 46-60 yrs, IV: 60 yrs and above), but the majority of informants (~35%) were above 60 years of age. These local people come from various ethnic cultures and poor educational backgrounds.

Table 2 — Informants, age group, and education level considered for documentation of traditional knowledge

Informants		Male	Female	Total
		Total number	81	65
	Percentage	55.48%	44.52%	
Age group	15-30 yrs	7	12	19 (13.01%)
	31-45 yrs	24	17	41 (28.08%)
	46-60 yrs	19	16	35 (23.97%)
	Above 60 yrs	31	20	51 (34.93%)
Education level	Illiterate	24	29	53 (36.30%)
	Attended school upto 5 classes	11	13	24 (16.44%)
	Attended school for 6-10 classes	22	11	33 (22.60%)
	Intermediate (12 th class) onwards	24	12	36 (24.66%)

Large proportions of the informants (~36%) were illiterate, while around ~64% of people had a secondary or a college level of education (Table 2).

Life-Forms, habit, habitat, and traditional uses

The ethnobotanical surveys revealed 28 indigenous Himalayan orchid species used as folklore medicine to be widely distributed in six Himalayan Indian states and one union territory, viz., Jammu and Kashmir, Himachal Pradesh, Uttarakhand, Sikkim, Arunachal Pradesh, Meghalaya, and Assam. These orchid plants are familiar with nomadic tribal people such as *Gujjars*, *Bakarwal*, *Himachaali*, *Garos*, *Khasis*, *Assammes*, *Arunchalis*, *Bhotiyas*, and *Jaunsari* residing in and around the hilly/mountainous regions of the study areas, and frequently employed in local preparation of ethnomedicine. These orchid species includes *Acampe ochracea* (Lindl.) Hochr., *Aerides odorata* Lour., *A. multiflora* Roxb., *Anoectochilus setaceus* Blume, *Arundina graminifolia* (D.Don) Hochr., *Bulbophyllum odoratissimum* (Sm.)

Table 3 — Morphology, part(s) used, traditional therapeutic uses, phytochemicals and biological properties of Himalayan orchids

Botanical name/Voucher number	Life form	Mode of Use	Part(s) used	Citation	Frequency (%)	Traditional therapeutic uses	Active chemical constituents	Biological properties
<i>Acampe ochracea</i> (Lindl.) Hochr. /116699	annual	paste	Lv	12	16.90	Fresh paste used in the treatment of the bone fracture.	praemorsin	Antimicrobial efficacy against kanamycin and ampicillin resistant strains of <i>E. coli</i> ²⁰ .
<i>Aerides odorata</i> Lour. / 118277	perennial	powder	Wp	32	45.07	Dried powder used in the treatment of tuberculosis.	ascorbic acid, apigenin, catechin, <i>p</i> -coumaric acid, ferulic acid, gallic acid, kaempferol, myricetin, naringin, quercetin, sinapic acid, syringic acid, rutin	Antimicrobial efficacy against kanamycin and ampicillin resistant strains of <i>E. coli</i> ²⁰ .
<i>Aerides multiflora</i> Roxb. /59292	perennial	paste	Tb, St	17	23.94	Paste used to heal wounds, boils and skin infections.	apigenin, catechin, <i>p</i> -coumaric acid, gallic acid, <i>p</i> -hydroxy, myricetin, rutin	Antibacterial properties ²⁰ .
<i>Anoectochilus setaceus</i> Blume / 50095	annual	powder	Wp	19	26.76	Dried powdered used as herbal tea, and decoction used in treatment of fever, pleurodynia, snake bite, lung infection, and hypertension.	flavonoids	Antimicrobial properties; <i>Xylaria</i> , isolated from leaves has helvolic acid, which is active against <i>Bacillus subtilis</i> and methicillin-resistant to <i>Staphylococcus aureus</i> ²⁴ .
<i>Arundina graminifolia</i> (D.Don.) Hochr. /64516	perennial	paste/ extract	Rz, Lv, St	4	5.63	Paste used to cure wounds and cuts; fresh juice obtained from young leaves and stems used in treatment of ear troubles.	arundinan, brundinaol, arundinoside A-G, arundiquinone & arundigramin rac-syringaresinol, astragalin, batatasin III, coelonin, densiflorol B, ephemeranthequinone, flavanthrin, gramniphinol H, lusian-thridin, orchinol, quercetin	polyphenols and flavonoids show antioxidant activity; Gramniphinol H shows cytotoxicity property against five tumor cell lines ^{2,20} .
<i>Bulbophyllum odoratissimum</i> (Sm.) Lindl. ex Wall. /116613	perennial	paste	Wp	12	16.90	Paste used in treatment of tuberculosis, chronic inflammation and fractured bones.	batatasin III, bulbobiphenantrene-bulbophythrins A and B coelonin, densiflorol B, gigantol, moscatin, 7-hydroxy-2,3,4-trimethoxy-9,10-dihydrophenanthrene, 1-monolinolein, syringaldehyde, syringing, tristin	Anticancerous; bulbobiphenantrene shows inhibition on human leukemia cell HL-60 and K562, human lung cancer cell strain A549 and human gastric carcinoma SGC-7901; densiflorol B and bulbobiphenantrene used as anticancerous drugs ²⁴ . data deficient.
<i>Coelogyne corymbosa</i> Lindl. /116786	annual	paste	Pb, St	12	16.90	Paste of pseudobulbs used on burns; stem paste applied on forehead to get relieve from headache.	coeloginone	

(Contd.)

Table 3 — Morphology, traditional therapeutic uses, phytochemicals and biological properties of Himalayan orchids (*Contd.*)

Botanical name/Voucher number	Life form	Mode of Use	Part(s) used	Citation	Frequency (%)	Traditional therapeutic uses	Active chemical constituents	Biological properties
<i>Coelogyne punctulata</i> Lindl. /60130	perennial	extract	Wp	9	12.68	Juice extract used to make vomits in case of poison.	coeloginone, coelogin, quabain, batatasin III	Data deficient.
<i>Crepidium acuminatum</i> (D.Don) Szlach. /64415	annual	extract/ decoction	Pb	14	19.72	Decoction used to cure fever, tuberculosis, bleeding diathesis and phthisis.	flavonoids, caryophyllene, eugenole, gallic acid, humulene, linoleic acid	Antioxidant activity against a long lived 2,2-diphenyl-1-picrylhydrazyl (DPPH) radical ³² .
<i>Cymbidium aloifolium</i> (L.) Sw. / 116709	perennial	paste/ extract	Wp	36	50.70	Roots paste plastered tightly to cure bones fractures; seeds useful in healing wounds and boils; leaves juice dropped in ears to cure otitis.	batatasin III, aloifol II, coumarin, cymbinodin A & B, methoxycoelonin.	Antibacterial; ethanol extracts show antinociceptive and anti-inflammatory activity against carrageenin induced paw oedema in mice ⁴² .
<i>Cymbidium elegans</i> Lindl./ 116719	perennial	extract	Tb, Fl	12	16.90	Fresh extract obtained from tubers and flowers used for vomit in poison or toxic substances.	cymbinodin	Data deficient.
<i>Dendrobium densiflorum</i> Wall. /116701	perennial	paste	Lv	28	39.44	Paste used in treatment of bones fractures and considered as calcium supplying plant.	dendroflorin, gigantol, scopoletin, moscatilin, scoparone, densiflorol A & B, moscatin, confusarin, dendrodensiflorol, cyripedin	Antimicrobial, anti-inflammatory activities; exhibits <i>in vitro</i> ant-platelet aggregation activity ⁴⁸ .
<i>Dendrobium fimbriatum</i> Hook. /101135	perennial	paste/ extract	Wp	17	23.94	Leaves juice used in liver upsets and for curing nervous debility; paste used in setting bones fractures.	moscatilin, gigantol, defuscin, chrysotoxine, 2,5-phenanthrenediol, 9-10-dihydro-7-methoxy, fimbriadimerbibenzyls	Cytotoxic properties against five human cell lines with IC ₅₀ value ranges from 2.2 to 21.2 µM.
<i>Dendrobium fugax</i> Rchb.f. /118587	perennial	powder	Tb	12	16.90	Taken as aphrodisiac, and used to cure insomnia.	salutaridine, thebaine	Data deficient.
<i>Dendrobium nobile</i> Lindl./28416	perennial	powder	Se	34	47.89	Grinded power applied on cuts, boils and wounds for quick healing; also taken in case of dry mouth and menstrual pains.	gigantol, denbinobine, dendramine, copacamphane, picrotoxane, cyclococamphane, nobiline, dendrophenol	Antioxidant activity; highly effective against Sarcoma 180 <i>in vivo</i> and HL-60 <i>in vitro</i> , showing antitumorous activity; denbinobine shows anticancer activity in human myelogenous K562 leukemia cells ²² .
<i>Eulophia graminea</i> Lindl. /53021	annual	extract	Rz	16	22.54	Juice extract used as tonic, and effective in curing cough and paralysis.	linalool, ferulic acid	Data deficient.

(Contd.)

Table 3 — Morphology, traditional therapeutic uses, phytochemicals and biological properties of Himalayan orchids (*Contd.*)

Botanical name/Voucher number	Life form	Mode of Use	Part(s) used	Citation	Frequency (%)	Traditional therapeutic uses	Active chemical constituents	Biological properties
<i>Habenaria dentata</i> (Sw.) Schltr. /63928b	annual	extract	Wp	29	40.85	Extract used in urinary problems; used as analgesic, disinfectant, aphrodisiac and antirheumatic.	habenariol, gastrodigenin	Apparently less explored; habenariol isolated used as anti-inflammatory agent ²² .
<i>Habenaria marginata</i> Colebr. /29138	annual	extract/ decoction	Tb	14	19.72	Decoction of fresh boiled tubers along with honey consumed daily in empty stomach to cure malignant ulcer.	diterpenes, hecogenin.	Data deficient.
<i>Herminium lanceum</i> (Thunb. ex Sw.) Vuijk /63928	annual	extract	Lv, St	7	9.86	Decoction prepared from green leaves and stems used to cure diabetes, fever and nose bleeding.	alkaloids	Data deficient.
<i>Liparis nervosa</i> (Thunb.) Lindl. /53324	annual	extract	Lv	37	52.11	Juice extract used in treatment of burns and ulcers.	linoleic acid, p-hydroxybenzaldehyde, swertisin, 6,8-di-C- α -L-arabinopyranosyl genkwanin, apigenin.	Data deficient.
<i>Papilionanthe teres</i> (Roxb.) Schltr. /42661	perennial	paste	Lv. St	10	14.08	Paste applied on forehead to keep brain cool during malaria fever, and also taken to protect body from cold and cough.	butanenediol, vandateroside I, II, III	Data deficient.
<i>Phaius tankervilleae</i> (Banks ex L'Her.) Blume / 61429	perennial	extract	Pb	13	18.31	Paste used to cure bones fractures, and extract taken to get relief from dysentery.	alkaloids, flavonoids	Data deficient.
<i>Pholidota imbricata</i> Lindl. /59456	perennial	extract	Tb	11	15.49	Paste used in treatment of stomach and rheumatic pains.	phoimbrtol A, loddigesiinol B, shanciol B, quercetin, luteolin, platycaryanin D, imbricatin.	Data deficient.
<i>Pholidota pallida</i> Lindl. / 116891	perennial	extract	Rz	8	11.27	Paste of fresh part used to cure finger abscess.	loddigesiinol B, luteolin, phenanthrene, phoimbrtol A, platycaryanin D, shanciol B.	Data deficient.
<i>Porpax muscicola</i> (Lindl.) Schuit., Y.P.Ng & H.A.Pedersen /53838	perennial	extract	Wp	33	46.48	Extract taken to get relief from chest, heart and lungs pain; juice of fresh leaves used to cure eye problems.	flavonoids, alkaloids	Data deficient
<i>Rhynchosyulis retusa</i> (L.) Blume /116697	perennial	extract	Rt, Lv	19	26.76	Fresh juices applied on cuts, boils and wounds; also taken orally in empty stomach to cure blood dysentery.	gigantol, flavidin,	Extract indicated positive antibacterial activity against <i>Pseudomonas aeruginosa</i> , <i>Staphylococcus aureus</i> , <i>E. coli</i> and <i>Salmonella typhi</i> ²⁰ .

(Contd.)

Table 3 — Morphology, traditional therapeutic uses, phytochemicals and biological properties of Himalayan orchids (*Contd.*)

Botanical name/Voucher number	Life form	Mode of Use	Part(s) used	Citation	Frequency (%)	Traditional therapeutic uses	Active chemical constituents	Biological properties
<i>Vanda coerulea</i> Griff. ex Lindl. /11872	perennial	extract	Lv	16	22.54	Fresh juice used in treatment of diarrhea and dysentery.	imbricatin, methoxycoelonin, gigantol, flavidin, coelonin, stilbenoids; stilbenoids (1-3) are the best stem biomarkers.	Hydro-alcoholic stem extract displayed DPPH/(•)OH radical scavenging activity ²⁰ .
<i>Vanda cristata</i> Wall. ex Lindl. /22523	perennial	extract	Lv	13	18.31	Fresh juice extracted used as a tonic and helps in body growth and to cure cough and cold.	gigantol, flavidin, coelonin	Data deficient.

(Lv-Leaves, Wp-Whole plant, St-Stem, Tb-Tuber, Rz-Rhizome, Pb- Pseudobulbs, Fl-Flower, Se-Seed)

Active phytochemicals constituents

Similar to other higher angiosperms, plant parts of orchid species do synthesize innumerable phytochemicals. Due to restrictions in a wide collection of orchid species, research on only a few species has been undertaken to investigate their biological properties and thousands of species are still unknown. Alkaloids, anthocyanins, carotenoids, flavonoids, and sterols are major groups of chemical constituents present in orchid species, and usually, alkaloids and flavonoids groups are most important for biological functions²⁰. The genus *Dendrobium* is a well-recognised genus studied for its phytochemical content and dendrobine was the first alkaloid investigated from Himalayan orchid *Dendrobium nobile*³⁸. *Dendrobium* is known to produce a variety of secondary metabolites in the form of alkaloids, bibenzyls, fluorenones, phenanthrens, and sesquiterpenes³², and these active phytochemical constituents are responsible for biological properties in human and animal health care. Several phenanthrene constituents (amoenumin, crepeditin, cumulatin dihydrophenanthrene), denfingenin, defuscin, dendrophenol, ephemerantherone, gigantol, moscatilin, moscatin, shihunidine, shihunine and rotundatin reported from orchids²¹⁻²³. The genus *Anoectochilus* Blume, *Arundina* Kunth, *Bulbophyllum* Thouars, *Bletilla* Rchb.f., *Cypripedium* L., *Eria* Lindl., *Eulophia* C. Agardh, *Gastrodia* R.Br., *Habenaria* Willd., *Malaxis* Sol. ex Sw., *Vanda* Jones ex R.Br. and *Vanilla* Mill. are enriched with different important phytochemicals³⁸. However, alkaloids present in orchids were studied by many researchers³⁹⁻⁴².

Lüning⁴³ studied the alkaloid content of 525 orchid species and recorded 214 species belonging to sixty

four genera contain 0.1% or more alkaloids. Approximately, 8% of *Dendrobium*, 18% of *Eria* and 42% of *Liparis* have this degree of alkaloid content. Zhang *et al.*⁴⁴ studied 42 species of *Dendrobium* having 100 compounds including 31 alkaloids, 6 coumarins, 15 bibenzyls, 4 fluorenones, 22 phenanthrenes, and 7 sesquiterpenoids. According to Hossain²⁰ more than 2000 species of orchid have been screened for their chemical constituents mostly rich in alkaloids and flavonoids contents. A study has shown that flavonol glycosides are mostly reported from Neottioid orchid species, whereas flavone glycosides are commonly found in the Epidendroid and Vandoid subfamily distributed in tropical and sub-tropical regions²⁰. The major active phytochemical constituents present in orchid species having a high demand in the pharmaceutical companies for medicine development are given in Fig. 2. Literatures^{2,39,45} screening showed pharmacological bioactivities for *A. multiflora*, *A. setaceus*, *A. grammifolia*, *B. odoratissimum*, *C. corymbosa*, *C. acuminatum*, *C. aloifolium*, *C. elegans*, *D. nobile*, *D. fimbriatum*, *E. graminea*, *F. fugax*, *H. marginata*, *H. lanceum*, *L. nervosa*, *P. teres*, *P. tankervilleae*, *P. imbricata*, *P. pallida*, *P. muscicola*, *R. retusa*, *V. Coerulea*, and *V. cristata*. This may be due to the shortage of planting materials for chemistry and pharmacological activities as most of the endemic orchids are categorized under the endangered category by IUCN, and wild collection from nature is strictly prohibited by State Forest Departments¹⁵.

The present investigation identifies the tribal applications of twenty-eight Himalayan orchids that occur in Himalayan belts. Chemical analyses indicate orchids are loaded with several commercially important bioactives in the form of alkaloids

Fig. 2 — Major bioactive phytochemicals of Himalayan orchids.

(dendrobine, jebantine), bibenzyls (gigantol, moscatilin), phenanthrenes (nudol, vllinol, agrostophyllinone, isoagrostophyllol, nidemin), 1-4 phenanthrenequinone (cypripedin), glycosides (loroglossin, melianin, kinsenoside), carbohydrates flavonoids, phenolic compounds (esters, glycosides) and stilbenoids (arundinan, orchinol, 6-methoxycoelonin, imbricatin, flaccidin, oxoflaccidin, isooxoflaccidin, flaccidin, agrostophyllin, callosin, callosinin, callosumin, callosuminin, callosumidin). Major commercially important compounds isolated and described from the Himalayan orchids includes agrostophyllin, agrostophyllinone, apigenin, arundinan, astragal, callosin, callosinin, callosumidin, callosumin, callosuminin, catechin, chrysotoxine, cypripedin, dendrobine, densiflorol B, ephemeranthoquinone, flaccidin, flaccidin, ferulic acid, gastrodigenin, gigantol, imbricatin,

isoagrostophyllol, isooxoflaccidin, jebantine, kaempferol, kinsenoside, loroglossin, luteolin, 6-methoxycoelonin, melianin, moscatilin, nidemin, nudol, orchinol, oxoflaccidin, phenanthrene, picrotoxane, quabain, quercetin, methyl gallate, myricetin, shanciol B, salutaridine, and thebaine. Many more compounds have been extracted from different parts (leaves, pseudobulbs, roots, stems, flowers, fruits) of the orchids⁴⁶⁻⁴⁸. *Dendrobium* and *Cymbidium* species have been used in traditional medicines in Taiwan, Korea, and Japan for the treatment of stomachache, healing fractures, kidney strengthening, fever, cancer treatment, and diabetic⁴⁹⁻⁵⁰. Several Ayurvedic medicines and nutraceutical products such as chyavanprasha, sudarshana churna, balarishtha, and amritprasha ghritam are available in the Indian markets whose formulations were prepared by using orchid species⁵¹⁻⁵³.

Conclusion

The ethnobiological investigations of the medicinal plant species of Himalayan regions and their uses among various tribal populations have proven that the medicinal potential of shared species were not dependent on geography, but depending on how frequently these species are used in the treatment of various diseases. Human-induced climate change has led to accelerated warming of the Himalayas in the recent past decade, which might have led to a decrease in the population of the biodiversity abode in different pockets of the regions. The orchids are a valuable source of bioactive compounds and are to be believed to possess medicinal properties that can help to treat many diseases. There is a need for inventory work for wild orchids from the conservation point of view, and this will help in the preservation of traditional knowledge associated with this group of plants, and for planning future research and developmental science associated with orchids or similar taxa.

Conflict of interest

Author declares no conflict of interest.

Acknowledgment

The author is thankful to the local people for their assistance in the field and for sharing their indigenous knowledge.

References

- Atwi M, Weiss E K, Loupassaki S, Makris D P, Ioannou E, *et al.*, Major antioxidant polyphenolic phytochemicals of three *Salvia* species endemic to the Island of Crete, *J Herbs Spices Med Plants*, 2016, **22**, 27–34.
- Bungtongdee N, Kathawut S, Laosripaiboon W and Lamtham S, The chemical composition, antifungal, antioxidant and antimutagenicity properties of bioactive compounds from fungal endophytes associated with Thai orchids, *J Phytopathol*, 2018, **167**(1), 56–64.
- Strobel G, Daisy B, Castillo U and Harper J, Natural products from endophytic microorganisms, *J Nat Prod*, 2004, **67**(2), 257–268.
- Musarella C M, Paglianiti I and Spampinato G, Ethnobotanical study in the poro and preserre Calabresi territory (Vibo Valentia, S-Italy), *Atti Soc Tosc Sci Nat Mem Ser B*, 2019, **126**, 13–28.
- Abdul M A, Ullah Z and Pieroni A, Wild food plant gathering among Kalasha, Yidgha, Nuristani and Khowar speakers in Chitral, NW Pakistan, *Sustainability*, 2020, **12**, 9176.
- Singh B, *Himalaya Orchids: Distribution and Taxonomy*, (Write and Print Publications, New Delhi, India), 2015.
- Jalal J S, Kumar P and Pangtey Y P S, Ethnomedicinal orchids of Uttarakhand, western Himalaya, *Ethnobot Leaflet*, 2008, **12**, 1227–1230.
- Nugraha A S, Triatmoko B, Wangchuk P and Keller P A, Vascular epiphytic medicinal plants as sources of therapeutic agents: Their ethnopharmacological uses, chemical composition, and biological activities, *Biomolecules*, 2020, **10**, 181.
- Shrestha M, Burd M, Garcia J E, Dorin A and Dyer A G, Colour evolution within orchids depends on whether the pollinator is a bee or a fly, *Plant Biol*, 2019, **21**(4), 745–752.
- Pritchard H W, Orchid seed storage: Historical perspective, current status, and future prospects for long term conservation, *Selbyana*, 1993, **14**, 89–104.
- Jalal J S, Status, threats and conservation strategies for orchids of western Himalaya, India, *J Threat Taxa*, 2012, **4**(15), 3401–3409.
- Yonzon R, Lama D, Bhujel R B and Rai S, Present availability status, diversity resources and distribution of medicinal orchid species in Darjeeling Himalaya of West Bengal, India, *Int J Pharm Nat Med*, 2013, **1**, 14–35.
- Singh B and Bedi Y S, Eating from raw wild plants in Himalaya: Traditional knowledge documentary on Sheena tribe along LoC Border in Kashmir, *Indian J Nat Prod Resour*, 2017, **8**(3), 269–275.
- Schmeller D S, Gruber B, Budrys E, Framsted E, Lengyel S, *et al.*, National responsibilities in European species conservation: A methodological review, *Conserv Biol*, 2008, **22**(3), 593–601.
- Wagensommer R P, Medagli P, Turco A and Perrino E V, IUCN Red list evaluation of the *Orchidaceae* endemic to Apulia Region (Italy) and considerations on the application of the IUCN protocol to rare species, *Nat Conserv Res*, 2020, **5**, 90–101.
- Singh B, Surmal O, Singh B, Singh S, Bhat M N, *et al.*, Folklore plants used in Tibetan mountain-based Sowa-Rigpa system of food and medicine: A close look on plant-people perception to herbal cure, In *Plants for Novel Drug Molecules-Ethnobotany to Ethnopharmacology*, edited by B Singh, Y P Sharma, (New India Publishing Agency, New Delhi, India), 2021, 1–43.
- Misra S, *Orchids of India-A Glimpse*, (Bishen Singh Mahendra Pal Singh, Dehradun, India), 2007.
- Sabin T P, Krishnan R, Vellore R, Priya P, Borgaonkar, *et al.*, Climate change over the Himalayas, In *Assessment of climate change over the Indian region*, edited by R Krishnan, J Sanjay, C Gnanaseelan, M Mujumdar, A Kulkarni, S Chakraborty, (Springer Nature, Switzerland), 2020.
- De L C and Medhi R P, Diversity and conservation of rare and endemic orchids of Northeast India -A review, *Indian J Hill Farm*, 2014, **27**(1), 81–89.
- Hossain M M, Therapeutic orchids: Traditional uses and recent advances - an overview, *Fitoterapia*, 2011, **82**, 102–140.
- Majumder P L and Pal S, Rotundatin, a new 9,10-dihydrophenanthrene derivative from *Dendrobium rotundatum*, *Phytochemistry*, 1992, **31**, 3225–3228.
- Zhang X, Xu J K, Wang J, Li W N, Kurihara H, *et al.*, Bioactive bibenzyl derivatives and fluorenones from *Dendrobium nobile*, *J Nat Prod*, 2007, **70**, 24–28.
- Li Y, Wang C L, Guo S X, Yang J S and Xiao P G, Two new compounds from *Dendrobium candidum*, *Chem Pharm Bull*, 2008, **56**, 1477–1479.

- 24 Singh S, Singh A K, Kumar S, Kumar M, Pandey P K, *et al.*, Medicinal properties and uses of orchids: A concise review, *Appl Bot*, 2012, **52**, 11627–11634.
- 25 Duthie J F, The orchids of the Northwestern Himalaya, *Ann Roy Bot Gard*, 1906, **9**(2), 81–211.
- 26 Raizada M B, Naithani H B and Saxena H O, *Orchids of Mussoorie*, (Bishan Singh Mahendra Pal Singh, Dehradun, India), 1981.
- 27 Jain S K and Mehrotra A, *A Preliminary Inventory of Orchidaceae in India*, (Publisher Botanical Survey of India, Howrah, India), 1984.
- 28 Kataki S K, *Orchids of Meghalaya*, (Government of Meghalaya, Shillong, India), 1986.
- 29 Sood S K, *Orchidaceae and Mankind*, (Publisher Deep Publications, New Delhi, India), 2006.
- 30 Nayar M P and Sastry A R K, *Red Data Book of Indian Plants*, vol. I–III, (Botanical Survey of India, Kolkata, India), 1987–1990.
- 31 Kumar C S and Manilal K S, *A Catalogue of Indian Orchids*, (Publisher Bishan Singh Mahendra Pal Singh, Dehradun, India), 1994.
- 32 Bose T K, Bhattacharjee S K, Das P and Basak U C, *Orchids of India*, (Publisher Naya Prakash, Calcutta, India), 1999.
- 33 Medhi R P and Chakrabarti S, Traditional knowledge of NE people on conservation of wild orchids, *Indian J Tradit Knowl*, 2009, **8**(1), 11–16.
- 34 Singh B and Borthakur S K, Wild medicinal plants used by tribal communities of Meghalaya, *J Econ Taxon Bot*, 2011, **35**, 331–339.
- 35 Singh B, Borthakur S K and Phukan S J, A survey of ethnomedicinal plants utilized by the indigenous people of Garo Hills with special reference to the Nokrek Biosphere Reserve (Meghalaya), India, *J Herbs Spices Med Plants*, 2014, **20**(1), 1–30.
- 36 Jain S K and Rao R R, *A Handbook of Field and Herbarium Methods*, (Today and Tomorrow's Printer Publisher, New Delhi, India), 1977.
- 37 Thiers B, Index Herbariorum: A global directory of public herbaria and associated staff, New York Botanical Garden's Virtual Herbarium, 2021, Available online: <http://sweetgum.nybg.org/science/ih>.
- 38 Bhattacharjee S K, *Herbaceous Perennials and Shade Loving Foliage Plants*, (Pointer Publishers Jaipur, India), 2006.
- 39 Arditti J, *Fundamentals of Orchid Biology*, (John Wiley & Sons, New York), 1992.
- 40 Bulpitt C J, The uses and misuses of orchids in medicine, *QJM Int J Med*, 2005, **98**(9), 625–631.
- 41 Chowdhery H J, *Orchid Flora of Arunachal Pradesh*, (Bishan Singh Mahendra Pal Singh, Dehradun, India), 1998.
- 42 Howlader M A, Mahmudul A, Ahmed K T, Khatun F and Sarker A A, Antinociceptive and anti-inflammatory activity of the ethanolic extract of *Cymbidium aloifolium* (L.) Sw., *Pak J Biol Sci*, 2011, **14**(19), 909–911.
- 43 Lüning B, Alkaloid content of orchidaceae, In *The orchids: scientific studies*, edited by C L Withner, (John Wiley and Sons, New York), 1974.
- 44 Zhang T X, Wang Y F and Cirello J, The herbal medicine tian ma gou teng alters the development of high blood pressure in the spontaneously hypertensive rat, *Am J Chin Med*, 1989, **17**, 211–219.
- 45 Brzosko E and Mirski P, Floral nectar chemistry in orchids: A short review and meta analysis, *Plants (Basel)*, 2021, **10**(11), 2315.
- 46 Sut S, Maggi F and Dall'Acqua S, Bioactive secondary metabolites from orchids (Orchidaceae), *Chem Biodivers*, 2017, **4**(11), e1700172.
- 47 Van-der-niet T, Jurgons A and Johnson S D, Pollinators, floral morphology and scent chemistry in the southern African orchid genus *Schizodutus*, *South Afr J Bot*, 2010, **76**(4), 726–738.
- 48 Ramya M, Jang S, An H R, Lee S Y, Park P M, *et al.*, Volatile organic compounds from orchids: From synthesis and function to gene regulation, *Int J Mol Sci*, 2020, **21**(3), 1160.
- 49 Fan C, Wang W, Wang Y, Qin G and Zhao W, Chemical constituents from *Dendrobium densiflorum*, *Phytochemistry*, 2001, **57**(8), 1255–1258.
- 50 Singh B, Singh B, Kishor A, Singh S, Bhat M N, *et al.*, Exploring plant-based ethnomedicine and quantitative ethnopharmacology: Medicinal plants utilized by the population of Jasrota Hill in Western Himalaya, *Sustainability*, 2020, **12**, 7526.
- 51 Singh B, *Plants for Human Survival and Medicine*, (CRC Press, Boca Raton, FL, USA; Taylor & Francis: London, UK), 2019.
- 52 Radhika B, Murthy J V V S N and Grace D N, Preliminary phytochemical analysis and antibacterial activity against clinical pathogens of medicinally important orchid *Cymbidium aloifolium* (L.) Sw., *Int J Pharm Sci Res*, 2013, **4**(10), 3925–3931.
- 53 Kong J M, Goh N K, Chia L S and Chia T F, Recent advances in traditional plant drugs and orchids, *Acta Pharmacol Sin*, 2003, **24**(1), 7–21.